

Discovery of a Nearby Low–Surface–Brightness Spiral Galaxy

W.B. Burton

University of Leiden Observatory, P.O. Box 9513, 2300 RA Leiden, Netherlands

R. Braun

Netherlands Foundation for Research in Astronomy, P.O. Box 2, 7990 AA Dwingeloo,
Netherlands

and

R.A.M. Walterbos and C.G. Hoopes

Department of Astronomy, New Mexico State University, Las Cruces, NM 88003

ABSTRACT

During the course of a search for compact, isolated gas clouds moving with anomalous velocities in or near our own Galaxy (Braun and Burton 1998), we have discovered, in the data of the Leiden/Dwingeloo survey (Hartmann and Burton 1997) of Galactic hydrogen, the H I signature of a large galaxy, moving at a recession velocity of 282 km s^{-1} with respect to our Galaxy. Deep multicolor and spectroscopic optical observations show the presence of star formation in scattered H II regions; radio H I synthesis interferometry confirms that the galaxy is rich in H I and has the rotation signature of a spiral galaxy; a submillimeter observation failed to detect the CO molecule. The radio and optical evidence combined suggest its classification as a low–surface–brightness spiral galaxy. It is located in close spatial and kinematic proximity to the galaxy NGC 6946. The newly–discovered galaxy, which we call Cepheus 1, is at a distance of about 6 Mpc. It is probably to be numbered amongst the nearest few LSB spirals.

Subject headings: galaxies: individual (Cepheus 1) — galaxies: ISM

1. Introduction

The census of even large, nearby galaxies is likely to be incomplete. Extinction of light due to intervening interstellar matter hides some 20% of the extragalactic sky

(Kraan–Korteweg and Lahav 1998, Balkowski and Kraan–Korteweg 1994, Seeberger, Saurer, and Weinberger 1996, Karachentseva and Karachentsev 1998) at optical wavelengths, prejudicing against the discovery of all types of galaxies. An additional bias pertains to the class of low–surface–brightness galaxies, which, although carrying substantial mass, have modest stellar luminosities because of their slow evolution. The bias against discovering LSB systems (Impey and Bothun 1997, Bothun, Impey, and McGaugh 1997, de Blok 1997) becomes even more severe under conditions of substantial extinction.

The Zone of Avoidance traces the accumulated optical extinction resulting from overlapping interstellar clouds, forming an irregular band along the Zodiac. The swath of obscuration is particularly broad in the constellation Cepheus. In 1920, Pannekoek had painstakingly drawn the contours of the Milky Way from counts of stellar surface density, and had devoted particular attention to the Cepheus region: his contours show a cleft of low stellar density (not recognized yet in 1920 as due to obscuration by dust) rising up from the Galactic equator and crossing the direction of the galaxy whose discovery is reported here. In 1934, Hubble had traced the distribution of some 44,000 extragalactic systems, and listed the region as the Cepheus Flare, which extends as far as 20° above the equator of the Milky Way, amongst those containing *no* galaxy.

The searches for hidden galaxies which rely on detecting the radio signals at 21-cm, specifically the Dwingeloo Obscured Galaxies Survey (Henning et al. 1998, Rivers et al. 1998) and the Parkes Multibeam Survey (Staveley–Smith et al. 1998), have been confined to about 5° of the Galactic equator and have given priority attention to directions where known cosmological structures, most notably the Supergalactic Plane, cross the Milky Way. The principal optical and infrared searches for hidden galaxies cover larger swaths of the sky, but did not detect this system of low optical brightness. The new galaxy is at a higher galactic latitude, 8° , than covered in radio ZoA searches, and is located at a large angle (some 45°) from the Supergalactic Plane defined by the galaxies in the Local Supercluster. The Nearby Galaxies Atlas (Tully and Fisher 1987), comprising 2367 galaxies with velocities less than 3000 km s^{-1} , shows only the galaxy NGC 6946 and its dwarf companion UGC 11583, within a circle of diameter more than 30° encompassing the Cepheus Flare region (except for NGC 6951 which, at a recession velocity of more than 1700 km s^{-1} , is so distant as to be irrelevant here). The obscuration in this region is substantial, but only about half that of the average at latitudes within 5° of the galactic equator. This so-called Local Void (Tully and Fisher 1987) is probably intrinsically sparsely populated, although, as we remark further below, the DOGS effort has found three probable additional galaxies accompanying NGC 6946.

2. Observations

2.1. Discovery as an Interloper amongst H I High-Velocity Clouds

Although substantially opaque at optical wavelengths, interstellar matter is largely transparent to radio radiation. The new galaxy was discovered during an effort (Braun and Burton 1998) to identify and characterize a class of compact, isolated high-velocity clouds. HVCs are objects detected in H I emission, moving at velocities inconsistent with the kinematics of the Milky Way as currently understood. Their distances remain largely uncertain (Wakker and van Woerden 1997); there is in particular no measure of the distance to any of the compact, isolated objects. The compact, broad H I signature of many of these objects resembles what one also might expect from a gas-rich nearby galaxy. H I investigation of such lines can be complicated by radio interference: we dealt with the possibility of interference contamination by using the Dwingeloo 25-meter telescope to observe, anew, all candidate members of the class of compact HVCs we had found in the Leiden/Dwingeloo survey (Hartmann and Burton 1997), on a dense, fully-sampled grid. This confirmed the reality of the signals (interference is not repeatable) and furthermore tightly constrained the direction of the unresolved features to an angular accuracy of about $5'$.

One spectrum amongst those of some 60 confirmed compact HVC's drew our particular attention. The spectrum shown in Fig. 1 resembles the discovery spectrum (Kraan-Korteweg et al. 1994) of the galaxy Dwingeloo 1, now recognized as the nearest, grand-design barred spiral: both show a broad, weak, H I signature at such a modest velocity as to blend with foreground emission from the Milky Way. We noticed also that the spectral feature in question had a velocity inconsistent with the other compact HVCs in our list. The recession velocity of the suspect feature is $v_{\text{LSR}} = 65 \text{ km s}^{-1}$ measured with respect to the Local Standard of Rest reference frame conventionally used in high-velocity cloud work; the corresponding heliocentric velocity is $v_{\text{HEL}} = 58 \text{ km s}^{-1}$. Correcting for the motion of the LSR within our Galaxy gives a velocity relative to the Galactic Standard of Rest of $v_{\text{GSR}} = v_{\text{LSR}} + 220 \cos(b) \sin(l) = 282 \text{ km s}^{-1}$. The kinematics of the class of compact HVCs (Braun and Burton 1998) resembles that of the members of the Local Group (Grebel 1997): but there is no Local Group member, and no other compact HVC, with a v_{GSR} as large as that measured for the suspect feature. We searched the standard catalogs of emission and reflection nebulae, planetary nebulae, and H II regions, finding no plausible alternative to an extragalactic system.

2.2. Optical Data

We extracted the field containing the suspect H I feature from the STScI Digitized Sky Survey CD-ROM, providing a scan of the optical Palomar Observatory Sky Survey red plate. Within the error circle determined by our new Dwingeloo data, we found a faint smudge, centered near α, δ (J2000) = $20^{\text{h}}51^{\text{m}}10^{\text{s}}.6, +56^{\circ}53'25''.0$, or $l, b = 94^{\circ}.37, +8^{\circ}.01$. We then contacted M. Irwin, who manages the APM microdensitometer at the Royal Greenwich Observatory. The scans he kindly produced gave higher resolution than the DSS, and, more importantly, included all three colors of the POSS II data; they confirmed the reality of the optical counterpart to the H I signal, and gave, in the color information, a discriminant against a Galactic reflection or emission nebula.

After the H I signature had been confirmed in new Dwingeloo data, and the optical counterpart found on the APM scans, we secured Director’s Discretionary Time on the ARC 3.5-meter telescope of the Apache Point Observatory, some Netherlands service time on the James Clerk Maxwell Telescope, for an attempt at a CO detection; and approval for detailed H I interferometry on the DRAO seven-antenna Synthesis Telescope in Penticton, British Columbia.

We observed the region with the 3.5-m telescope at Apache Point Observatory to determine the nature of the optical counterpart. First, using the DIS camera, green (Gunn g: 5240 Å) and red (Gunn r: 6700 Å) images were obtained of a single $4' \times 4'$ field centered on the POSS II optical smudge. Subsequently, using the SPICAM camera, we imaged a $7' \times 14'$ region in the broadband red (R: 6600 Å) and near-infrared (I: 8000 Å) bands, using the drift-scanning mode to obtain a larger field of view. The effective exposure time was 10 minutes in both filters; the image resolution was typically between $1''.2$ and $1''.5$. We also observed a single $4'.4 \times 14'$ strip for 5 minutes in a narrow (25 Å passband) filter centered at 6561 Å, close to the rest wavelength of the H α line. This allowed a search for ionized nebulae created by massive hot stars in the velocity range $-600 < v_{\text{LSR}} < +500 \text{ km s}^{-1}$. We removed continuum emission from stars in the narrow filter by subtracting a suitably-scaled R-band image. Some 30 H II regions were detected in this short exposure, distributed over a spatial extent of $5'.5$ diameter. Figure 2 shows a color composite of the optical data.

The detection of H II regions was important because optical spectroscopy of H II regions is a powerful tool to further elucidate the nature of Cepheus 1. We obtained spectra of two H II regions close to the optical center of the galaxy with the DIS spectrograph on the 3.5-m telescope, covering wavelengths from 4600 to 5300 Å, and from 5700 to 6800 Å. The radial velocities of the H II regions are $v_{\text{LSR}} = 50 \pm 5 \text{ km s}^{-1}$ and $v_{\text{LSR}} = 58 \pm 5 \text{ km s}^{-1}$, respectively. We derived the total optical extinction towards the H II regions from the flux ratio of the H α to H β recombination lines. The Balmer-decrement values are $A_V = 2.45$

and $A_V = 2.78$, respectively, in good agreement with the Galactic foreground extinction of $A_V = 2.0$ deduced from the Galactic H I column density, because H II regions typically have 0.5 to 1.0 magnitudes of extinction due to dust in the galaxy in which they are embedded. Taking this internal extinction into account, we adopt $A_V = 2.0$, consistent with the foreground H I column and the corrected Balmer decrements. (We note that extinction extrapolated from the DIRBE dust analysis by Schlegel, Finkbeiner, and Davis (1998) yields $A_V = 3.1$; our determination of the extinction using the Balmer decrement is a direct one, and refers to the exact direction of the galaxy, and thus is to be preferred over the extrapolated, relatively low resolution, DIRBE result.)

We determined the optical morphology and integrated stellar luminosity of Cepheus 1 from the broadband images, after correcting for Galactic foreground extinction. We summed the optical light in elliptical apertures, after removing foreground stars from the images. The light profile can be described by a single exponential with a scale length of $60''$ (1.8 kpc for the distance of 6 Mpc defended below), but there is considerable uncertainty in the exact shape because of the faintness of the emission and the large number of foreground stars. The best fitting ellipses are strongly flattened (axial ratio about 0.3 to 0.5), which seems in contradiction with the much more circular appearance of the H I and H II region distributions. It is possible that the continuum light we detect is mainly contributed by a faint stellar bulge or bar, while the actual disk may be mostly too faint to be seen at all; the H II regions extend well beyond the radius to which we can trace the continuum light. However, if we do attribute the light detected here to a disk, we derive the following parameters. The central disk surface brightness, corrected for foreground extinction, is $21.9 \text{ mag arcsec}^{-2}$ in R, and 22.8 to $23.0 \text{ mag arcsec}^{-2}$ in B. (Galaxies with such central surface brightness would be selected against, even under moderate foreground extinction.) Cepheus 1 has a faint stellar disk which, in the current data, can only be traced over an angular diameter of $3'$, about half the extent of the H II region distribution. Converting the de-reddened g–r color, derived from the calibrated DIS images, to standard Johnson passbands following Jorgensen (1994), we find $B-V=0.57$, typical for a disk galaxy.

2.3. Additional Radio Data

Radio observations of the H I emission line were carried out with the DRAO Synthesis Telescope, covering a 2° field over the velocity range $-340 < v_{\text{LSR}} < +500 \text{ km s}^{-1}$. The resolution was $69'' \times 60''$ NS by EW and 4 km s^{-1} in velocity. The resulting sensitivity in four twelve-hour tracks was 13.3 mJy/beam in each velocity channel. H I emission from Cepheus 1 extends at least over the v_{LSR} range $+7$ to $+133 \text{ km s}^{-1}$. The low-velocity

emission is somewhat confused with H I emission from the Galaxy, so the total velocity extent may still be underestimated. However, the systemic (LSR) velocity of the galaxy is well defined by the observed kinematics described below to be $65 \pm 1 \text{ km s}^{-1}$. Using the positive-velocity flank of the H I emission profile and the systemic velocity allows determination of the linewidths at 20% and 50% of the peak intensity, $W_{20} = 124 \text{ km s}^{-1}$ and $W_{50} = 90 \text{ km s}^{-1}$. An image of the integrated H I column depth is shown in Figure 3; the corresponding velocity field is shown in Figure 4. Figure 3 shows a high-brightness gaseous disk with a diameter of about $7'$ in which column densities between 5×10^{20} to 10^{21} cm^{-2} are observed. The H I diameter defined by an azimuthally averaged face-on surface density of $1 \text{ M}_{\odot} \text{ pc}^{-2}$ is $11'.7$ and the integrated H I flux is 136 Jy km s^{-1} . The small ellipticity of the high-brightness H I disk is consistent with inclinations of less than about 35° . Over this region, the kinematics are well described by a rotating disk system with constant inclination and a position angle of the receding line of nodes of about -50° . The kinematics indicate a spiral galaxy. The rotation velocity in the inner disk rises relatively slowly to a maximum value which is closely linked to the assumed inclination, but this parameter is difficult to determine, from the current data, for such a closely face-on system. If the inclination is as large as 35° the rotation velocity would only be about 60 km s^{-1} , but for an inclination of 20° , it would be about 100 km s^{-1} , and it would be even larger for a smaller inclination.

Beyond the bright, inner H I disk, there is a discontinuous change in the kinematics. An offset in the iso-velocity contours of some 20 km s^{-1} is observed, in the sense of a shift to higher apparent rotation velocity. For this reason, the most extreme velocities are found in the outer-disk H I arms that extend NW and S of the galaxy center. The outer-disk kinematics can be modeled by an increasing inclination and shifting of the line of nodes, but this solution is poorly constrained by the limited position angles at which outer disk gas is detected. A less likely possibility, although one which can not yet be ruled out, is that the inclination stays fixed while the rotation velocity continues to climb. If this were the case, the rotation velocity would climb to 100 km s^{-1} or 190 km s^{-1} , for inclinations of 35° or 20° , respectively.

The observed parameters of Cepheus 1 are summarized in Table 1.

3. Physical Properties of Cepheus 1

Having established the presence of Cepheus 1, we considered its distance and physical properties. Random and systematic motions of nearby galaxies can be at least as large as

those expected from the Hubble flow, rendering the cosmological velocity/distance relation of little use for systems nearer than about 10 Mpc. A preliminary distance estimate follows from the tendency of galaxies to cluster on all scales. A galaxy is not commonly more remote from any neighbor than, say, 1 Mpc. The case of NGC 6946, one of the 3 or 4 nearest giant Sc spiral galaxies, is interesting in this regard. It occupies an apparently lonely position (at $l, b = 95^\circ.7, 11^\circ.7$) in the Local Void (Tully and Fisher 1987), accompanied by its dwarf companion UGC 11583 at a projected distance of $38'$, corresponding approximately to the linear distance of the Large Magellanic Cloud from our own Galaxy. The distance of NGC 6946 has recently been determined (Sharina, Karachentsev, and Tikhonov 1997) as 6.4 ± 0.4 Mpc, from photometry of resolved stars in it and its irregular dwarf companion, UGC 11583, using the value $A_V = 1.93$ to correct for the foreground extinction. The heliocentric velocity of NGC 6946 is $+51 \pm 4$ km s $^{-1}$, corresponding to $v_{\text{GSR}} = 280$ km s $^{-1}$.

In view of its angular and, especially, kinematic, proximity to NGC 6946, it seems altogether plausible that Cepheus 1 is at a similar distance. Cepheus 1 is situated south of NGC 6946 some $3^\circ.9$, corresponding at 6 Mpc to a projected linear distance of 410 kpc, which is about half the distance separating the Milky Way from our own nearest large neighbor, the Andromeda galaxy. The kinematic separation between NGC 6946 and the new galaxy is only 2 km s $^{-1}$ in the v_{GSR} frame. Recently the Dwingeloo Obscured Galaxies Survey (Henning et al. 1998, Rivers et al. 1998) detected 21-cm signatures from three candidate galaxies within about 10° of NGC 6946 and each with a recessional velocity $v_{\text{LSR}} < 250$ km s $^{-1}$. If, as seems likely, these candidates are confirmed in optical and radio synthesis data to be dwarf galaxies, then NGC 6946 and its accompanying systems will constitute a small group.

The distance of 6 Mpc, suggested as a working hypothesis, can be tested in several ways. The APO H α image shows H II regions scattered over the entire field; a deeper image of a larger field would plausibly reveal more such regions. If we adapt $6'$ as the angular dimension and 6 Mpc as the distance, then the projected linear size of the H II region distribution of Cepheus 1 is 10.5 kpc. The linear size following from the $11'.7$ H I dimension is 21 kpc; the total H I gas mass is $M_{\text{HI}} = 1.1 \times 10^9 M_\odot$. Cepheus 1 is thus comparable to the nearby spiral M33 (Corbelli, Schneider, and Salpeter 1989, Guidoni, Messi, and Natali 1981) in terms of the scale length of the stellar continuum (1.7 kpc for M33), the spatial extent of the H II region distribution (15 kpc diameter for M33), and the total H I mass ($1 \times 10^9 M_\odot$). The H I diameter of Cepheus 1 (21 kpc) is larger than that of M33 (15 kpc when defined in the same way). The disk of M33 is somewhat bluer ($B-V=0.46$) than that of Cepheus 1, and intrinsically brighter, by a factor 6 in total brightness and a factor 3 or 4 in central surface brightness. Of the 19 LSB galaxies in the sample of de Blok (1997), 7 have an H I size smaller than that of Cepheus 1, and 12 have a larger size. Of the 48

HSB galaxies in Broeils’s (1992) “Sample 1”, 13 have an H I size smaller than Cepheus 1, and 35 have a larger H I size. Galaxies with an H I diameter of about 20 kpc are typically classified as Sc or Sd. Comparisons with the properties of other nearby LSB systems (see Karachentseva et al. 1996), in particular with those of nearest member of that class, NGC 247 (Carignan 1985), will be made when additional data are available. We note that the H I diameter of Cepheus 1 is about as large as that measured for the Milky Way to the distance at which the H I surface density of our own Galaxy begins to rapidly fall.

We determined the integrated magnitudes of Cepheus 1 from the APO data. The apparent magnitude in the R band, integrating the disk profile to infinity, is in the range $m_R = 13.1$ to 13.5 . Correcting for foreground extinction this implies $m_R = 11.5$ to 11.9 . The absolute magnitudes for $D = 6$ Mpc are in the range $M_R = -17.4$ to -17.0 and $M_B = -16.5$ to -16.1 . The faintness of its disk is consistent with Cepheus 1 being an LSB galaxy, albeit a rather bright example of these systems. The cautions expressed in Section 2.2 apply here as well, i.e. the optical disk parameters are not yet well constrained, which could affect the integrated optical magnitudes, and the actual central surface brightness estimate, which could be lower if there is in fact a faint bar present. The uncertainties stem from the range of possible disk scale lengths. No correction has yet been made for internal extinction within that disk, which may be low, if the dust-to-gas ratio is low; the H I column with a normal dust-to-gas ratio would give $\tau_V = 0.2 - 0.4$. Assuming an internal extinction in the B band of $A_B = 0.3$ mag, suggests an absolute extinction-corrected B magnitude of about -16.6 mag. The resulting ratio M_{HI}/L_B is $1.6 M_\odot/L_\odot$, for a value $M_B = 5.48$ for the Sun. This is a very high ratio of gas mass to optical luminosity for normal high-surface-brightness galaxies, and even moderately high for the class of LSB systems (de Blok 1997).

A further check on the distance follows from the luminosities of the H II regions. We derived the cumulative H α luminosity function of the H II regions in Cepheus 1, and compared it with that for M33. Taking foreground extinction into account, the two luminosity functions overlap if the distance to Cepheus 1 is taken as 6 Mpc. While this constraint is not very strong, given the possible variation in H II –region luminosity functions, a distance very different than 6 Mpc seems unlikely. We have not yet derived a Tully–Fisher distance to the new galaxy. In view of the intrinsic scatter (Zwaan et al. 1995) in the T–F relationship for LSB systems, and in view of the uncertainties in the inclination, i.e. in Δv , we postpone this derivation until the more detailed data expected are available.

The optical spectra also support the LSB classification. Notably, we only detected emission from doubly-ionized oxygen in the H II regions, but not from singly-ionized nitrogen or sulphur. For the brighter H II region, we derived limits on the line ratios of

$[\text{NII}](6584 \text{ \AA})/\text{H}\alpha < 0.10$ and $[\text{SII}](6716 \text{ \AA})/\text{H}\alpha < 0.12$, and a high ratio of $[\text{OIII}](5007 \text{ \AA})/\text{H}\beta = 2.8$. These line ratios are entirely consistent with typical LSB H II regions (Impey and Bothun 1997, Bothun, Impey, and McGaugh 1997), but not with the inner H II regions in normal spirals. The ratios indicate a low abundance of heavy elements, a characteristic property of LSB galaxies. An attempt was made on the JCMT to detect CO(J=2–1) from the central direction of Cepheus 1. That the attempt returned a null spectrum is not decisive, because spiral galaxies show a wide range of molecular morphologies relative to H I morphologies, but the lack of detectable CO is consistent with the situation (de Blok 1997) in all other LSB systems. Because of the proximity of Cepheus 1, a very deep CO observation would plausibly establish a very good limit to the molecular content of an LSB galaxy. We note also the sparsely populated environment, where the evolution of Cepheus 1 proceeds with evidently little external influence on the internal physics.

In view of the various uncertainties, our distance determination should be regarded as preliminary, although the suggested association with NGC 6946 at about 6 Mpc is supported by all the data at our disposal. The evidence points toward the morphology of a low-surface-brightness spiral galaxy: the faint disk surface brightness, the scattered H II regions, the substantial H I columns, the relatively strong and regular H I rotation signature, the weak molecular level, the optical color, the low heavy-metal abundance, and the relative isolation in the Local Void. Cepheus 1 is one of the nearest few LSB spirals and the nearest seen at low inclination. It is a galaxy of moderate H I size, and a rather extreme $M_{\text{HI}}/L_{\text{B}}$ ratio. The discovery of nearby galaxies extends the knowledge of the local luminosity and mass functions. The discovery of a likely member of the LSB class offers opportunities for studies in detail. Such opportunities are rare because luminous LSB galaxies are mostly found at large distances, but exploiting what opportunities there are is important in view of the contributions such systems may make to understanding the formation and evolution of galaxies. By itself, Cepheus 1 would not significantly distort the dynamics of the Local Group. But it will be interesting to establish if NGC 6946 and Cepheus 1 together with the other gas-rich members which are being found (Henning et al. 1998, Rivers et al. 1998) nearby might, as a group, influence the gravitational torques in the local neighborhood (see Peebles 1994 and Lynden-Bell 1981). A multi-wavelength observational campaign has been prepared which will more tightly constrain the distance to the newly-unveiled galaxy and provide details of its structure.

We are grateful to M. Irwin for scanning the three-color POSS II data using the RGO APM machine; to E. Turner for rapidly providing access to the APO 3.5-meter telescope; to A. Gray and T. Landecker at the DRAO for facilitating H I mapping using the Penticton array; and to R. Tilanus and F. Baas for the CO observation at the JCMT.

REFERENCES

- Balkowski, C., and Kraan–Korteweg, R.C. 1994, Unveiling large-scale structures behind the Milky Way. ASP Conf. Ser., 67
- Bothun, G., Impey, C., and McGaugh, S. 1997, PASP, 109, 745
- Braun, R., and Burton, W.B. 1998, A&A, in press
- Broeils, A.H. 1992, Dark and visible matter in spiral galaxies, Ph.D. Thesis, University of Groningen
- Carignan, C. 1985, ApJS, 58, 107
- Corbelli, E., Schneider, S.E., and Salpeter, E.E. 1989, AJ, 97, 390
- de Blok, W.J.G. 1997, The properties and evolution of low surface brightness galaxies, Ph.D. Thesis, University of Groningen
- Grebel, E.K. 1997, Rev. in Modern Astr. 10, 29
- Guidoni, U., Messi, R., and Natali, G. 1981, A&A, 96, 215
- Hartmann, D., and Burton, W.B. 1997, Atlas of Galactic Neutral Hydrogen, Cambridge: Cambridge University Press
- Henning, P.A., Kraan–Korteweg, R.C., Rivers, A.J., Loan, A.J., Lahav, O., and Burton, W.B. 1998, AJ, 115, 584
- Hubble, E. 1934, ApJ, 79, 8
- Impey, C., and Bothun, G. 1997, ARA&A, 35, 267
- Jorgensen, I. 1994, PASP, 106, 967
- Karachentseva, V.E., and Karachentsev, I.D. 1998, A&AS, 127, 409
- Karachentseva, V.E., et al. 1996, A&AS, 117, 343
- Kraan–Korteweg, R.C., and Lahav, O. 1998, Scientific American, October issue
- Kraan–Korteweg, R.C., Loan, A.J., Burton, W.B., Lahav, O., Ferguson, H.C., Henning, P.A., and Lynden-Bell, D. 1994, Nature, 372, 77
- Lynden–Bell, D. 1981, Observatory, 101, 111

- Pannekoek, A. 1920, Die Nördliche Milchstrasse, Annalen Sterrewacht Leiden, XI, Derde Stuk
- Peebles, P.J.E. 1994, ApJ, 429, 43
- Rivers, A.J., Henning, P.A., Kraan–Korteweg, R.C., Lahav, O., Burton, W.B. 1998, Proc. Cardiff Workshop, in press
- Schlegel, D.J., Finkbeiner, D.P., and Davis, M. 1998, ApJ, 500, 525
- Seeberger, R., Saurer, W., and Weinberger, R. 1996, A&AS, 117, 1
- Sharina, M.E., Karachentsev, I.D., and Tikhonov, N.A. 1997, Astr. Letters, 23, 373
- Staveley–Smith, L., Juraszek, S., Koribalski, B.S., Ekers, R.D., Green, A.J., Haynes, R.F., Henning, P.A., Kesteven, M.J., Kraan–Korteweg, R.C., Price, R.M., and Sadler, E.M. 1998, AJ, in press
- Tully, R.B., and Fisher, J.R. 1987, Nearby Galaxies Atlas, Cambridge: Cambridge University Press
- Wakker, B.P., and van Woerden, H. 1997, ARA&A, 35, 217
- Zwaan, M.A., van der Hulst, J.M., de Blok, W.J.G., and McGaugh, S.S. 1995, MNRAS, 273, L35

Fig. 1.— H I spectrum of the galaxy Cepheus 1 extracted from the Leiden/Dwingeloo survey (Hartmann and Burton 1997), toward $l, b = 94^{\circ}.5, 8^{\circ}.0$. The velocity is relative to the Local Standard of Rest, the intensity is in terms of brightness temperature. The lower-velocity wing of the galaxy’s signature is buried under emission from our own Galaxy; thus from this detection spectrum alone it is not possible to determine the total velocity width of the galaxy’s signature, nor the velocity centroid giving the systemic velocity.

Fig. 2.— Color-coded composite of the I-, R-, and H α images of Cepheus 1 observed with the 3.5-m telescope of the Apache Point Observatory. The field size is $4'.4 \times 6'.8$ EW by NS. The red, green, and blue intensities in the image are proportional to the surface brightness in the I-band, R-band, and continuum-subtracted H α exposures, respectively. Note the bar-like concentration of stellar continuum emission peaking near α, δ (J2000) = $20^{\text{h}}51^{\text{m}}10^{\text{s}}.6, +56^{\circ}53'25''.0$, and the H II regions which are visible rather uniformly scattered over a large portion of the field. H α spectra were obtained for two H II regions, E and W of the center.

Fig. 3.— Integrated H I emission in Cepheus 1 as observed with the DRAO Synthesis Telescope. The linear pseudo-color scale extends over apparent column densities of 100 to $800 \times 10^{18} \text{ cm}^{-2}$ observed with a beam size of $69'' \times 60''$ NS by EW. The high-brightness inner disk appears nearly circular, suggesting an almost face-on orientation. With a size of about $7'$, the inner disk extends over a comparable field to that depicted optically in Figure 2. Fainter arm-like features extend further to both the North and South for a total gaseous extent of some $14'$. The observed N_{HI} corresponds to a hydrogen mass of $1.1 \times 10^9 M_{\odot}$, at a distance of 6 Mpc.

Fig. 4.— Velocity field of the H I emission in Cepheus 1 as observed with the DRAO Synthesis Telescope. The linear pseudo-color scale extends between $v_{\text{LSR}} = +10$ and $+100 \text{ km s}^{-1}$. Note the classic spider-diagram pattern within the inner disk, signifying the well-ordered rotation characteristic of a spiral galaxy. The outer disk shows a kinematic discontinuity of more than 20 km s^{-1} with respect to the adjacent inner disk velocities, suggesting either strong warping or a rising rotation curve in the outer disk.

Table 1. Properties of Cepheus 1

Parameter	Value	Notes
Optical center: $\alpha, \delta(\text{J2000})$	$20^{\text{h}}51^{\text{m}}10^{\text{s}}.6(\pm 5''), +56^{\circ}53'25''.0(\pm 5'')$	
Dynamical center: $\alpha, \delta(\text{J2000})$	$20^{\text{h}}51^{\text{m}}10^{\text{s}}.8(\pm 5''), +56^{\circ}53'26''.4(\pm 5'')$	
Galactic coordinates: l, b	$94^{\circ}375, +8^{\circ}010$	
Distance	6 Mpc	^a
Optical axis ratio	0.4 ± 0.1	^b
Foreground extinction, A_V	< 2.45 mag	^c
Adopted foreground extinction, A_V	2.0 ± 0.2 mag	^d
Internal extinction, A_B	0.3 ± 0.2 mag	^e
Central R surface brightness, μ_0^{R}	21.9 ± 0.2 mag arcsec ⁻²	^{b,f}
Central B surface brightness, μ_0^{B}	22.9 ± 0.2 mag arcsec ⁻²	^{b,f}
Absolute R magnitude, M_{R}	-17.5 ± 0.3 mag	^g
Absolute B magnitude, M_{B}	-16.6 ± 0.3 mag	^g
Total blue luminosity, L_{B}	$6.8 \pm 2.2 \times 10^8 L_{\text{B}\odot}$	^{g,h}
Stellar disk scalelength, s_R	60 arcsec, 1.8 kpc	^b
Total H I flux, F_{HI}	136 ± 10 Jy km s ⁻¹	
Total H I mass, M_{HI}	$1.1 \pm 0.1 \times 10^9 M_{\odot}$	
Systemic velocity, v_{HEL}	$+58 \pm 3$ km s ⁻¹	
Line width at 50% intensity, W_{50}	90 ± 5 km s ⁻¹	
Line width at 20% intensity, W_{20}	124 ± 5 km s ⁻¹	
Inclination	$28 \pm 8^{\circ}$	
Position angle	$-50 \pm 3^{\circ}$	
Diameter of H I disk, D_{HI}	11'7, 21 kpc	ⁱ

^aAssuming association with NGC 6946 as discussed in the text.

^bMay be strongly affected by the presence of a bar.

^cBased on Balmer decrements of two H II regions.

^dBased on Galactic N_{HI} .

^eBased on Cepheus 1 N_{HI} .

^fCorrected for foreground extinction.

^gCorrected for foreground and internal extinction.

^hAssuming $M_{\text{B}\odot} = 5.48$.

ⁱMeasured at a mass surface density of $1 M_{\odot} \text{ pc}^{-2}$.

This figure "fig1.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810432v1>

This figure "fig2.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810432v1>

This figure "fig3.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810432v1>

This figure "fig4.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810432v1>